

Quantumness tests and witnesses in the tomographic–probability representation

S. N. Filippov^{1*} and V. I. Man'ko^{2†}

¹ *Moscow Institute of Physics and Technology (State University)
Institutskii per. 9, Dolgoprudnyi, Moscow Region 141700, Russia*

² *P. N. Lebedev Physical Institute, Russian Academy of Sciences
Leninskii Prospect 53, Moscow 119991, Russia*

In view of the tomographic–probability representation of quantum states, we reconsider the approach to quantumness tests of a single system developed in [Alicki and Van Ryn 2008 *J. Phys. A: Math. Theor.* **41** 062001]. For qubits we introduce a general family of quantumness witnesses which are operators depending on an extra parameter. Spin tomogram and dual spin tomographic symbols are used to study qubit examples and the test inequalities which are shown to satisfy simple relations within the framework of the standard probability theory.

PACS numbers: 03.65.Ta, 03.65.Sq, 03.65.Wj

Keywords: quantumness witness, quantumness test, tomographic–probability representation of quantum states, spin tomogram

I. INTRODUCTION

A boundary between quantum and classical worlds is rather difficult to draw precisely, while the problem of their distinguishing is getting more important than it was hardly ever before. This problem is particularly important for the large-scale quantum computer to be constructed because it must be a macroscopic object and exhibit quantum properties at the same time. The investigations in this field of science are also stimulated by the attempts to describe quantum mechanical phenomena by different kinds of hidden variables models. Though these problems have been paid a great attention for many years the common point of view has not been achieved. The discussions came up with a bang after a recent proposal [1, 2, 3, 4] of a simple test of checking whether it is possible or not to describe a given set of experimental data by a classical probabilistic model.

In quantum mechanics the state of a system can be identified with a fair probability called tomographic probability distribution or state tomogram (see, e.g., the review [5]). The probability representation of quantum states with continuous variables (position, momentum) was introduced in [6] and that of states with discrete variables (spin, qudits) was introduced in [7, 8]. In the probability representation of quantum mechanics the relation between classical and quantum systems behavior can be studied using the same notion of states expressed in terms of the probability distribution (tomogram) in both classical and quantum domains [9, 10]. The quantum (or classical) nature of a system state can be clar-

*Electronic address: filippovsn@gmail.com

†Electronic address: manko@sci.lebedev.ru

ified if in the system the uncertainty relations [11, 12, 13, 14] for conjugate variables are fulfilled. Also, if the state is quantum, it can be similar to a classical one and there are some studies of the classicality (see, e.g., [15]) based on properties of the diagonal representation of density operator [16] (or P -function [17]).

In the works [1, 2], quantum or classical properties of a system state were associated with measuring some specific observables such that there exist certain inequalities which hold true in classical domain and are violated in quantum domain. Violation of the inequalities is considered as a quantumness witness of the system state. In this sense, the criterion [1, 2] is similar in its spirit to the Bell inequalities [18, 19]. The Bell inequalities were studied by means of the tomographic-probability representation in [20, 21, 22, 23]. The aim of our work is to consider the inequalities introduced in [1, 2] and their properties in classical and quantum domains within the framework of the probability representation of quantum states. We suppose that such a procedure is necessary while dealing with the quantum probabilistic model based on tomograms.

The paper is organized as follows.

In Sec. II, we are aimed at recalling the tomographic representation of qubit states and observables by employing both ordinary and dual tomographic symbols. In Sec. III, the quantumness test is discussed within the framework of classical and quantum probability descriptions. In Sec. IV, we present a family of observables which can be used to detect quantumness of an arbitrary generally mixed state $\hat{\rho} \neq \hat{1}/n$ of a single system. Here we also predict what kind of experiment one should make to test the quantumness of a state specified by its tomogram. In Sec. V, conclusions and prospects are presented.

II. STATES AND OBSERVABLES IN THE TOMOGRAPHIC-PROBABILITY REPRESENTATION

Apart from being described by the conventional density operator $\hat{\rho}$, the state of a qubit is determined thoroughly by its spin tomogram. The probability distribution function (spin tomogram) $w(m, u)$ is nothing else but the probability to obtain m ($m = \pm \frac{1}{2}$) as spin projection on the direction given by the unitary 2×2 matrix u . This matrix can be considered as a matrix of an irreducible representation of the rotation group depending on two Euler angles determining the direction of quantization (point on the Bloch sphere). The relation between $\hat{\rho}$ and $w(m, u)$ reads

$$w(\mathbf{x}) \equiv w(m, u) = \langle m | u \hat{\rho} u^\dagger | m \rangle = \text{Tr} \left(\hat{\rho} u^\dagger | m \rangle \langle m | u \right) = \text{Tr} \left(\hat{\rho} \hat{U}(\mathbf{x}) \right), \quad (1)$$

where the operator $\hat{U}(\mathbf{x}) = u^\dagger | m \rangle \langle m | u$ is called the dequantizer operator and $\mathbf{x}$ is assigned to the set of parameters (m, u) . The general procedure to use dequantizer operators was discussed in the context of star-product quantization schemes in [24, 25]. The explicit form of the dequantizer $\hat{U}(\mathbf{x})$ can be obtained readily by exploiting the matrix u expressed in terms of the Euler angles α, β, γ :

$$u = \begin{pmatrix} \cos(\beta/2) e^{-i(\alpha+\gamma)/2} & -\sin(\beta/2) e^{i(\alpha-\gamma)/2} \\ \sin(\beta/2) e^{-i(\alpha-\gamma)/2} & \cos(\beta/2) e^{i(\alpha+\gamma)/2} \end{pmatrix}. \quad (2)$$

Therefore, taking advantage of $|m\rangle\langle m| = \frac{1}{2}\hat{I} + m\hat{\sigma}_z$, where $\hat{I}$ is the 2×2 identity matrix and $\hat{\sigma}_z$ is the third Pauli matrix, one can write

$$\hat{U}(\mathbf{x}) = \frac{1}{2}\hat{I} + m\hat{F}(\alpha, \beta), \quad (3)$$

where $\mathbf{x} = (m, \alpha, \beta, \gamma)$ and the matrix $\hat{F}(\alpha, \beta)$ has the following form

$$\hat{F}(\alpha, \beta) = \begin{pmatrix} \cos \beta & -e^{i\alpha} \sin \beta \\ -e^{-i\alpha} \sin \beta & -\cos \beta \end{pmatrix}. \quad (4)$$

If given the spin tomogram $w(\mathbf{x})$ it is possible to reconstruct the density operator $\hat{\rho}$ [26]. This reconstruction was shown to take the simple form

$$\hat{\rho} = \int w(\mathbf{x}) \hat{D}(\mathbf{x}) d\mathbf{x}, \quad (5)$$

where the integration implies

$$\int d\mathbf{x} = \sum_{m=-1/2}^{1/2} \int_0^{2\pi} d\alpha \int_0^\pi \sin \beta d\beta \int_0^{2\pi} d\gamma, \quad (6)$$

and the quantizer operator $\hat{D}(\mathbf{x})$ is defined by the formula

$$\hat{D}(\mathbf{x}) = \frac{1}{8\pi^2} \left(\frac{1}{2}\hat{I} + 3m\hat{F}(\alpha, \beta) \right). \quad (7)$$

In quantum mechanics any observable A is identified with a Hermitian operator $\hat{A}$. By analogy with the density operator $\hat{\rho}$ one can introduce the tomographic symbol $w_A(\mathbf{x})$ of the operator $\hat{A}$. Just in the same way we write

$$w_A(\mathbf{x}) = \text{Tr} \left(\hat{A} \hat{U}(\mathbf{x}) \right), \quad \hat{A} = \int w_A(\mathbf{x}) \hat{D}(\mathbf{x}) d\mathbf{x}. \quad (8)$$

It is worth noting that both quantizer and dequantizer are operators depending on the set of parameters $\mathbf{x}$ so it seems possible to swap quantizer with the dequantizer. Substituting the quantizer operator for the dequantizer one and visa versa leads to a so-called dual tomographic symbol $w_A^d(\mathbf{x})$ [25, 27] satisfying the following relations:

$$w_A^d(\mathbf{x}) = \text{Tr} \left(\hat{A} \hat{D}(\mathbf{x}) \right), \quad \hat{A} = \int w_A^d(\mathbf{x}) \hat{U}(\mathbf{x}) d\mathbf{x}. \quad (9)$$

The dual symbol in the tomographic-probability representation turned out to provide the function introduced in [9] which after averaging with a tomogram yields the mean value of the observable. Let us now express the average value of the observable A by means of ordinary and dual tomographic symbols. Indeed, the mean value of A reads

$$\text{Tr} (\hat{\rho} \hat{A}) = \text{Tr} \int d\mathbf{x} w(\mathbf{x}) \hat{D}(\mathbf{x}) \hat{A} = \int d\mathbf{x} w(\mathbf{x}) \text{Tr} (\hat{A} \hat{D}(\mathbf{x})) = \int w(\mathbf{x}) w_A^d(\mathbf{x}) d\mathbf{x}. \quad (10)$$

The formula obtained can be checked immediately for a general case of the density operator $\hat{\rho}$ and the observable A :

$$\hat{\rho} = \begin{pmatrix} \rho_{11} & \rho_{12} e^{i\zeta} \\ \rho_{12} e^{-i\zeta} & \rho_{22} \end{pmatrix}, \quad \hat{A} = \begin{pmatrix} A_{11} & A_{12} e^{i\eta} \\ A_{12} e^{-i\eta} & A_{22} \end{pmatrix}, \quad (11)$$

where ρ_{ij} , A_{ij} , ζ , η are real numbers. Then the spin tomogram $w(\mathbf{x})$ is

$$w(\mathbf{x}) = \frac{1}{2} + m(\rho_{11} - \rho_{22}) \cos \beta - 2m\rho_{12} \cos(\zeta - \alpha) \sin \beta \quad (12)$$

and the dual tomographic symbol $w_A^d(\mathbf{x})$ reads

$$w_A^d(\mathbf{x}) = \frac{1}{8\pi^2} \left[\frac{1}{2}(A_{11} + A_{22}) + 3m(A_{11} - A_{22}) \cos \beta - 6mA_{12} \cos(\eta - \alpha) \sin \beta \right]. \quad (13)$$

The direct calculation yields

$$\int w(\mathbf{x}) w_A^d(\mathbf{x}) d\mathbf{x} = \frac{1}{2}(A_{11} + A_{22}) + \frac{1}{2}(\rho_{11} - \rho_{22})(A_{11} - A_{22}) + 2\rho_{12}A_{12} \cos(\zeta - \eta) \quad (14)$$

that coincides totally with the quantity $\text{Tr}(\hat{\rho}\hat{A})$ computed by using the matrix form of operators (11).

Let us now express the average value of the observable A in terms of its possible outcomes $A_\uparrow$ and $A_\downarrow$ measured through separate experiments. The numbers $A_\uparrow$ and $A_\downarrow$ are nothing else but eigenvalues of the operator $\hat{A}$. Consequently there exists a unitary matrix u_A such that the matrix $\hat{A}$ can be factorized as follows

$$\hat{A} = u_A^\dagger \begin{pmatrix} A_\uparrow & 0 \\ 0 & A_\downarrow \end{pmatrix} u_A. \quad (15)$$

The matrix $u_A^\dagger$ is composed of two columns which are eigenvectors of the operator $\hat{A}$ corresponding to eigenvalues $A_\uparrow$ and $A_\downarrow$, respectively. In order to specify the matrix u_A one can substitute the Euler angles (ϕ, θ, φ) for (α, β, γ) in formula (2), i.e., $u_A = u(\phi, \theta, \varphi)$. Then the dual tomographic symbol of the operator $\hat{A}$ takes the form

$$w_A^d(\mathbf{x}) = \frac{1}{8\pi^2} \left\{ \frac{1}{2}(A_\uparrow + A_\downarrow) + 3m(A_\uparrow - A_\downarrow) \times [\cos \beta \cos \theta + \cos(\alpha - \phi) \sin \beta \sin \theta] \right\}. \quad (16)$$

Now, when $w_A^d(\mathbf{x})$ is known, it is not impossible to evaluate the integral

$$\begin{aligned} \int w(\mathbf{x}) w_A^d(\mathbf{x}) d\mathbf{x} &= \frac{1}{2}(A_\uparrow + A_\downarrow) + \frac{1}{2}(A_\uparrow - A_\downarrow)(\rho_{11} - \rho_{22}) \cos \theta - (A_\uparrow - A_\downarrow)\rho_{12} \cos(\zeta - \phi) \\ &= A_\uparrow \left[\frac{1}{2} + \frac{1}{2}(\rho_{11} - \rho_{22}) \cos \theta - \rho_{12} \sin \theta \cos(\zeta - \phi) \right] \\ &\quad + A_\downarrow \left[\frac{1}{2} - \frac{1}{2}(\rho_{11} - \rho_{22}) \cos \theta + \rho_{12} \sin \theta \cos(\zeta - \phi) \right] \\ &= w(+\frac{1}{2}, \phi, \theta, \varphi) A_\uparrow + w(-\frac{1}{2}, \phi, \theta, \varphi) A_\downarrow \\ &= w_\uparrow(u_A) A_\uparrow + w_\downarrow(u_A) A_\downarrow, \end{aligned} \quad (17)$$

that gives the average value of the observable A , i.e., the value of quantity $\text{Tr}(\hat{\rho}\hat{A})$. Here we denoted $w(m = +\frac{1}{2}, u)$ and $w(m = -\frac{1}{2}, u)$ by $w_\uparrow(u)$ and $w_\downarrow(u)$, respectively. One cannot help mentioning that the same result is achieved by using the definition of the spin tomogram (1):

$$\begin{aligned}
\text{Tr}(\hat{\rho}\hat{A}) &= \text{Tr}\left[\hat{\rho}u_A^\dagger\begin{pmatrix} A_\uparrow & 0 \\ 0 & A_\downarrow \end{pmatrix}u_A\right] = \text{Tr}\left[u_A\hat{\rho}u_A^\dagger\begin{pmatrix} A_\uparrow & 0 \\ 0 & A_\downarrow \end{pmatrix}\right] \\
&= \text{Tr}\left[\begin{pmatrix} \langle\frac{1}{2}|u_A\hat{\rho}u_A^\dagger|\frac{1}{2}\rangle & \langle\frac{1}{2}|u_A\hat{\rho}u_A^\dagger|-\frac{1}{2}\rangle \\ \langle-\frac{1}{2}|u_A\hat{\rho}u_A^\dagger|\frac{1}{2}\rangle & \langle-\frac{1}{2}|u_A\hat{\rho}u_A^\dagger|-\frac{1}{2}\rangle \end{pmatrix}\begin{pmatrix} A_\uparrow & 0 \\ 0 & A_\downarrow \end{pmatrix}\right] \\
&= \langle\frac{1}{2}|u_A\hat{\rho}u_A^\dagger|\frac{1}{2}\rangle A_\uparrow + \langle-\frac{1}{2}|u_A\hat{\rho}u_A^\dagger|-\frac{1}{2}\rangle A_\downarrow \\
&= w_\uparrow(u_A)A_\uparrow + w_\downarrow(u_A)A_\downarrow.
\end{aligned} \tag{18}$$

III. QUANTUMNESS TESTS

In this section, we are going to answer the question whether it is possible or not to describe the system involved by a classical probabilistic model. The negative reply indicates straightway the quantumness of the system in question. To start, let us consider the case of a single qubit, and then discuss a generalization for an arbitrary qudit system.

In the classical statistical model of a single qubit system the observable A is associated with a set of real numbers $(A_\uparrow, A_\downarrow)$, where the numbers $A_\uparrow$ and $A_\downarrow$ are nothing else but possible outcomes of the measurement of A . Moreover, the observable A^2 has possible outcomes of the form $(A_\uparrow^2, A_\downarrow^2)$, i.e., is in strong relation to A . The states form a simplex of probability distributions $(p_\uparrow, p_\downarrow)$, with $0 \leq p_\uparrow$, $0 \leq p_\downarrow$, and $p_\uparrow + p_\downarrow = 1$ (such a simplex is a geometrical treatment of all possible classical states; geometric interpretation of quantum states is reviewed in [28]). Then the expectation values of observables A and A^2 read

$$\langle A \rangle_{cl} = p_\uparrow A_\uparrow + p_\downarrow A_\downarrow, \quad \langle A^2 \rangle_{cl} = p_\uparrow A_\uparrow^2 + p_\downarrow A_\downarrow^2. \tag{19}$$

It is worth noting that such a classical system is equivalent to the quantum one, with both the density operator and the operator $\hat{A}_{cl}$ being of the diagonal form

$$\hat{\rho}_{cl} = \begin{pmatrix} p_\uparrow & 0 \\ 0 & p_\downarrow \end{pmatrix}, \quad \hat{A}_{cl} = \begin{pmatrix} A_\uparrow & 0 \\ 0 & A_\downarrow \end{pmatrix}. \tag{20}$$

Suppose one has two nonnegative observables A and B such that inequality $\langle A \rangle_{cl} \leq \langle B \rangle_{cl}$ holds true for all states $(p_\uparrow, p_\downarrow)$. That implies the following relations: $0 \leq A_\uparrow \leq B_\uparrow$ and $0 \leq A_\downarrow \leq B_\downarrow$. If this is the case, the average value of observable A^2 is necessarily less or equal than the average value of B^2 . Indeed,

$$\langle A^2 \rangle_{cl} = p_\uparrow A_\uparrow^2 + p_\downarrow A_\downarrow^2 \leq p_\uparrow B_\uparrow^2 + p_\downarrow B_\downarrow^2 = \langle B^2 \rangle_{cl}. \tag{21}$$

The mathematical aspect of the classical statistical model is briefly expressed by the implication

$$\text{Tr}\left[\begin{pmatrix} p_\uparrow & p_\uparrow \\ p_\downarrow & p_\downarrow \end{pmatrix}\begin{pmatrix} A_\uparrow & A_\downarrow \\ -B_\uparrow & -B_\downarrow \end{pmatrix}\right] \leq 0 \quad \Rightarrow \quad \text{Tr}\left[\begin{pmatrix} p_\uparrow & p_\uparrow \\ p_\downarrow & p_\downarrow \end{pmatrix}\begin{pmatrix} A_\uparrow^2 & A_\downarrow^2 \\ -B_\uparrow^2 & -B_\downarrow^2 \end{pmatrix}\right] \leq 0, \tag{22}$$

where we introduced a stochastic matrix with the matrix elements $p_\uparrow$ and $p_\downarrow$.

In the quantum statistical model the observable A is associated with the operator $\hat{A}$, which in its turn corresponds to the Hermitian matrix whose eigenvalues $A_\uparrow$ and $A_\downarrow$ give

possible outcomes of the measurement of A . States are identified with density operators $\hat{\rho}$, which are positive Hermitian matrices with the trace equal to unity. By using the tomographic representation of quantum states and observables developed in previous section, one can write the average values of observables A and A^2 as follows:

$$\langle A \rangle_q = \text{Tr}(\hat{\rho}\hat{A}) = w_{\uparrow}(u_A)A_{\uparrow} + w_{\downarrow}(u_A)A_{\downarrow}, \quad (23)$$

$$\langle A^2 \rangle_q = \text{Tr}(\hat{\rho}\hat{A}^2) = w_{\uparrow}(u_A)A_{\uparrow}^2 + w_{\downarrow}(u_A)A_{\downarrow}^2. \quad (24)$$

In a way analogous to the classical case, we consider two nonnegative operators $\hat{A}$ and $\hat{B}$ (i.e., having nonnegative eigenvalues $A_{\uparrow}, A_{\downarrow}, B_{\uparrow}, B_{\downarrow} \geq 0$) such that the residual operator $\hat{B} - \hat{A}$ is nonnegative as well. The last requirement can be rewritten in the form of the condition

$$\text{Tr} \left[\begin{pmatrix} w_{\uparrow}(u_A) & w_{\uparrow}(u_B) \\ w_{\downarrow}(u_A) & w_{\downarrow}(u_B) \end{pmatrix} \begin{pmatrix} A_{\uparrow} & A_{\downarrow} \\ -B_{\uparrow} & -B_{\downarrow} \end{pmatrix} \right] \leq 0 \quad (25)$$

that must be valid for *all* tomograms $w(m, u)$. In (25), we introduced a stochastic matrix with the matrix elements $w_{\uparrow}(u_{A,B})$ and $w_{\downarrow}(u_{A,B})$. In contrast to the classical case, even under these circumstances the average value of the operator $\hat{B}^2 - \hat{A}^2$ does not have to be nonnegative. Therefore, there can exist the tomogram $\tilde{w}(m, u)$ such that

$$\text{Tr} \left[\begin{pmatrix} \tilde{w}_{\uparrow}(u_A) & \tilde{w}_{\uparrow}(u_B) \\ \tilde{w}_{\downarrow}(u_A) & \tilde{w}_{\downarrow}(u_B) \end{pmatrix} \begin{pmatrix} A_{\uparrow}^2 & A_{\downarrow}^2 \\ -B_{\uparrow}^2 & -B_{\downarrow}^2 \end{pmatrix} \right] > 0. \quad (26)$$

The difference between classical and quantum behaviour of the observable $A^2 - B^2$ serves as the basis for a simple quantumness test.

Quantumness Test. Two observables A and B are found which averaged values satisfy the inequality $0 \leq \langle A \rangle \leq \langle B \rangle$ for all experimentally accessible states, but for a certain state the condition $\langle A^2 \rangle \leq \langle B^2 \rangle$ is violated, then the state involved cannot be described by a classical probabilistic model and is surely to be quantum.

This is the reason why the operator $\hat{B}^2 - \hat{A}^2$ was proposed to call quantumness witness [2].

The comparison of the expressions (22) and (26) shows that despite both classical and quantum cases can be treated probabilistically the difference between them occurs due to the classical state being associated with numbers $(p_{\uparrow}, p_{\downarrow})$ while the quantum one being identified with the functions $w_{\uparrow}(u)$ and $w_{\downarrow}(u)$ taking on the different values at the unitary group elements $u = u_A$ and $u = u_B$.

The generalization of this approach for a qudit state given by its $n \times n$ density matrix $\hat{\rho} \neq \hat{1}/n$ is evident while dealing with operators $\hat{A}$, $\hat{B}$ having an effect on the two-dimensional subspace spanned by eigenvectors corresponding to two different eigenvalues of $\hat{\rho}$.

IV. QUANTUMNESS WITNESSES

The quantumness test described above requires a certain quantumness witness to be found. In the earlier work [2], the abundance of quantumness witnesses has been demonstrated for qubit states $\hat{\rho} \neq \frac{1}{2}\hat{I}$. In spite of that, the proposed scheme does not give the explicit shape of the operators $\hat{A}$ and $\hat{B}$. In this section, we are going to repair this gap.




FIG. 1: The average value $\text{Tr} [\hat{\rho}_d(\hat{B}_d^2 - \hat{A}_d^2)]$ (expectation value of quantumness witness) is negative for all r of the range $0 < r \leq 1$ ($a = 0.35$, $b = 0.1$, and the values of k are depicted in the figure).

First of all, the state $\hat{\rho} = \frac{1}{2}\hat{I}$ can be treated as classical, because its tomogram $w(m, u) \equiv \frac{1}{2}$ and coincides with the case of classical probabilities $p_{\uparrow} = p_{\downarrow} = \frac{1}{2}$. Let us demonstrate that all the other qubit states are quantum.

Indeed, an arbitrary density operator $\hat{\rho}$ can be reduced to the diagonal form of

$$\rho_d = \begin{pmatrix} \frac{1+r}{2} & 0 \\ 0 & \frac{1-r}{2} \end{pmatrix} \quad (27)$$

that becomes maximally mixed (classical) if $r = 0$ and pure if $r = 1$. In order to find the quantumness witness for all $0 < r \leq 1$ we avoid limitations $\text{Tr}\hat{B} = 1$ or $\text{Tr}\hat{B} = 2$ imposed in [1] and [2], respectively. Below we introduce a family of the operators $\hat{A}$ and $\hat{B}$ that satisfy the requirements $0 \leq \hat{A} \leq \hat{B}$ while $\hat{B}^2 - \hat{A}^2$ is a quantumness witness for a particular region of parameters a, b, k . For the diagonal density matrix (27) one can use

$$\hat{A}_d = \begin{pmatrix} \frac{1}{2r^2} - \frac{a}{r} & \frac{1}{r}\sqrt{\frac{1}{4r^2} - a^2} \\ \frac{1}{r}\sqrt{\frac{1}{4r^2} - a^2} & \frac{1}{2r^2} + \frac{a}{r} \end{pmatrix}, \quad (28)$$

$$\hat{B}_d = \hat{A}_d + \begin{pmatrix} br^k & -br^k \\ -br^k & br^k \end{pmatrix}. \quad (29)$$

The operators $\hat{A}_d$, $\hat{B}_d$, $\hat{B}_d - \hat{A}_d$ have nonnegative eigenvalues only while the average value $\text{Tr} [\hat{\rho}_d(\hat{B}_d^2 - \hat{A}_d^2)]$ is negative for all r of the range $0 < r \leq 1$. Figure 1 gives the illustration of this fact for particular values of parameters a, b, k .

All-tomographic statement of the quantumness test can be formulated as follows.

Suppose we have an experimentally measured spin tomogram $w(m, u)$ of a qubit state. The parameter r of the diagonal density matrix $\hat{\rho}_d$ can be found as

$$r = \max_{u \in SU(2)} (w_{\uparrow}(u) - w_{\downarrow}(u)) = w_{\uparrow}(u_{\rho}) - w_{\downarrow}(u_{\rho}). \quad (30)$$

Then for the operators $\hat{A}$ and $\hat{B}$ leading to the quantumness witness one can use

$$\hat{A} = u_{\rho}^{\dagger} \hat{A}_d(r) u_{\rho}, \quad \hat{B} = u_{\rho}^{\dagger} \hat{B}_d(r) u_{\rho}. \quad (31)$$

V. CONCLUSIONS

To conclude we summarize the main results of the paper. We demonstrated that the criterion of quantumness of a system state found in [1, 2] can be formulated within the framework of the probability representation of quantum states. In this representation, the structure of the criterion is clarified since it is formulated by means of inequalities, where stochastic matrices involved (see (22) and (26)) have different properties. In case of classical states the stochastic matrix providing inequality has constant matrix elements. Conversely, in case of quantum states the stochastic matrix providing analogous inequality has matrix elements depending on unitary group elements. In view of this, for some values of the unitary group elements the inequality can be violated. One can extend the analysis of quantumness tests given for qubits to systems with continuous variables, e.g., for photon quadrature components.

Acknowledgments

V. I. M. thanks the Russian Foundation for Basic Research for partial support under Project Nos. 07-02-00598 and 08-02-90300.

-
- [1] Alicki R and Van Ryn N 2008 *J. Phys. A: Math. Theor.* **41** 062001
 - [2] Alicki R, Piani M and Van Ryn N 2008 *J. Phys. A: Math. Theor.* **41** 495303
 - [3] Alicki R 2007 Quantumness of Josephson junctions reexamined *Preprint* quant-ph/0610008
 - [4] Alicki R 2008 On von Neumann and Bell theorems applied to quantumness tests *Preprint* quant-ph/0810.4708
 - [5] Man'ko V I, Marmo G, Simoni A, Sudarshan E C G and Ventrigilia F 2008 *Rep. Math. Phys.* **61** 337
 - [6] Mancini S, Man'ko V I and Tombesi P 1996 *Phys. Lett. A* **213** 1
 - [7] Dodonov V V and Man'ko V I 1997 *Phys. Lett. A* **229** 335
 - [8] Man'ko V I and Man'ko O V 1997 *J. Exp. Ther. Phys.* **85** 430
 - [9] Man'ko O and Man'ko V I 1997 *J. Russ. Laser Res.* **18** 407
 - [10] Man'ko V I and Mendes R V 2000 *Physica D* **145** 330
 - [11] Heisenberg W 1927 *Z. Phys.* **43** 172
 - [12] Schrödinger E 1930 *Ber. Kgl. Akad. Wiss. Berlin* **24** 296
 - [13] Robertson H P 1930 *Phys. Rev.* **35** 667
 - [14] Robertson H P 1934 *Phys. Rev.* **46** 794
 - [15] Shchukin E V and Vogel W 2005 *Phys. Rev. A* **72** 043808

- [16] Sudarshan E C G 1963 *Phys. Rev. Lett.* **10** 277
- [17] Glauber R J 1963 *Phys. Rev.* **131** 2766
- [18] Bell J S 1964 *Physics* **1** 195
- [19] Clauser J F, Horne M A, Shimony A and Holt R A 1969 *Phys. Rev. Lett.* **23** 880
- [20] Lupo C, Man'ko V I and Marmo G 2007 *J. Phys. A: Math. Theor.* **40** 13091
- [21] Lupo C, Man'ko V I and Marmo G 2006 *J. Phys. A: Math. Gen.* **39** 12515
- [22] Andreev V A, Man'ko V I, Man'ko O V and Shchukin E V 2006 *Theor. Math. Phys.* **146** 172
- [23] Filippov S N and Man'ko V I 2009 *J. Russ. Laser Res.* **30** 49
- [24] Man'ko O V, Man'ko V I and Marmo G 2002 *J. Phys. A: Math. Gen.* **35** 699
- [25] Man'ko O V, Man'ko V I, Marmo G and Vitale P 2007 *Phys. Lett. A* **360** 522
- [26] Castaños O, López-Peña R, Man'ko M A and Man'ko V I 2003 *J. Phys. A: Math. Gen.* **36** 4677
- [27] Man'ko V I, Marmo G and Vitale P 2005 *Phys. Lett. A* **334** 1
- [28] Filippov S N and Man'ko V I 2008 *J. Russ. Laser Res.* **29** 564